

for demonstration of the method, we note from Table 4 that 95% of the population represented by the NHANES III data was likely to be exposed to ≤ 67 ppb chloroform in water, which can be compared with the MCL for trihalomethanes of 100 ppb (U.S. EPA 2007). Using distributions for the posterior concentration of chloroform in tap water (C_W), daily water intake (W_I), and body weight (BW), we drew 10,000 samples from the product $C_W W_I / BW$ to obtain a daily chloroform intake from drinking water (Table 5). The RfD for chloroform is 0.01 mg/kg/day (U.S. EPA 2001). The 95th percentile for the posterior distribution for chloroform concentration in ambient air is 0.02 $\mu\text{g}/\text{L}$. The U.S. EPA currently does not have an established inhalation reference concentration for chloroform (U.S. EPA 2001).

The accuracy of the results is limited by the approximate nature of the model, the assumptions regarding exposure, and the quality of the experimental data. In particular, the prior distribution from the TEAM data did not correspond to the same population as that of the NHANES III data, and although it provided an informative prior, data from the same locations and time frame as the NHANES III data, even if not corresponding to the individuals in that study, would likely improve the accuracy of the results. Although the method presented here is intended to be a tool to reconstruct exposure from biomonitoring data where no corresponding exposure data are available, comparison of the results with such data would greatly assist in assessing the accuracy of the

method, and such results could be incorporated as prior distributions for additional chloroform dose reconstructions.

CORRECTION

In the abstract of the original manuscript published online, the units for chloroform exposures in tap water were presented as milligrams per liter instead of micrograms per liter. They have been corrected here.

REFERENCES

- Allen BC, Hack CE, Clewell HJ. 2007. Use of Markov chain Monte Carlo analysis with a physiologically-based pharmacokinetic model of methylmercury to estimate exposures in US women of childbearing age. *Risk Anal* 27(4):947–959.
- Bernillon P, Bois FY. 2000. Statistical issues in toxicokinetic modeling: a Bayesian perspective. *Environ Health Perspect* 108(suppl 5):883–893.
- Bois FY, Maszle DR. 1997. MCSim: a Monte Carlo simulation program. *J Stat Soft* 2:1–60.
- Brown RP, Delp MD, Lindstedt SL, Rhomberg LR, Beliles RP. 1997. Physiological parameter values for physiologically based pharmacokinetic models. *Toxicol Ind Health* 13(4):407–484.
- Carlin BP, Louis TA. 2000. Bayes and Empirical Bayes Methods for Data Analysis. Boca Raton, FL:Chapman and Hall/CRC.
- Centers for Disease Control and Prevention. 1996. Third National Health and Nutrition Examination Survey, III (NHANES III) Public-Use Data Files. Available: http://www.cdc.gov/nchs/products/elec_prods/subject/nhanes3.htm [accessed 7 July 2008].
- Corley RA, Gordon SM, Wallace LA. 2000. Physiologically based pharmacokinetic modeling of the temperature-dependent dermal absorption of chloroform by humans following bath water exposures. *Toxicol Sci* 53(1):13–23.
- Covington TR, Robinan Gentry P, Van Landingham CB, Andersen ME, Kester JE, Clewell HJ. 2007. The use of Markov chain Monte Carlo uncertainty analysis to support a Public Health Goal for perchloroethylene. *Regul Toxicol Pharmacol* 47(1):1–18.
- Gelman A, Bois F, Jiang JM. 1996. Physiological pharmacokinetic analysis using population modeling and informative prior distributions. *J Am Stat Assoc* 91(436):1400–1412.
- Gelman A, Carlin JB, Stern HS, Rubin DB. 2004. Bayesian Data Analysis. Boca Raton, FL:Chapman and Hall/CRC Press.
- Gelman A, Rubin DB. 1996. Markov chain Monte Carlo methods in biostatistics. *Stat Methods Med Res* 5(4):339–355.
- Georgopoulos PG, Roy A, Gallo MA. 1994. Reconstruction of short-term multiroute exposure to volatile organic-compounds using physiologically-based pharmacokinetic models. *J Expo Anal Environ Epidemiol* 4(3):309–328.
- Gilks WR, Spiegelhalter DJ, Richardson SS. 1996. Markov Chain Monte Carlo in Practice. Boca Raton, FL:Chapman and Hall/CRC Press.
- Gnuplot. 2007. Gnuplot 4.0. Available: <http://www.gnuplot.info/> [accessed 21 October 2007].
- Hack CE, Chiu WA, Jay Zhao Q, Clewell HJ. 2006. Bayesian population analysis of a harmonized physiologically based pharmacokinetic model of trichloroethylene and its metabolites. *Regul Toxicol Pharmacol* 46(1):63–83.
- Hadarnard J. 1902. Sur les problemes aux derivees partielles et leur signification physique. *Bull Univ Princeton* 13:49–52.
- Hastings WK. 1970. Monte-Carlo sampling methods using Markov chains and their applications. *Biometrika* 57(1):97–106.
- Hindmarsh A. 1983. ODEPACK, a systemized collection of ODE solvers. In: *Scientific Computing* (Stepleman RS, ed). Amsterdam:North-Holland, 55–64.
- Jonsson F. 2001. Physiologically Based Pharmacokinetic Modeling in Risk Assessment: Development of Bayesian Population Methods. Uppsala, Sweden:Uppsala University.
- Marino DJ, Clewell HJ, Gentry PR, Covington TR, Hack CE, David RM. 2006. Revised assessment of cancer risk to dichloromethane: part I Bayesian PBPK and dose-response modeling in mice. *Regul Toxicol Pharmacol* 45(1):44–54.
- Metropolis N, Rosenbluth AW, Rosenbluth MN, Teller AH, Teller E. 1953. Equation of state calculations by fast computing machines. *J Chem Phys* 21(6):1087–1092.
- Miller G, Martz HF, Little TT, Guilmette R. 2002. Bayesian internal dosimetry calculations using Markov chain Monte Carlo. *Radiat Prot Dosimetry* 98(2):191–198.
- National Research Council. 2006. Human Biomonitoring for Environmental Chemicals. Washington, DC:National Academies Press.
- Roy A, Weisel CP, Gallo MA, Georgopoulos PG. 1996. Studies of multiroute exposure/dose reconstruction using physiologically based pharmacokinetic models. *Toxicol Ind Health* 12(2):153–163.
- Sohn MD, McKone TE, Blancato JN. 2004. Reconstructing population exposures from dose biomarkers: inhalation of trichloroethylene (TCE) as a case study. *J Expo Anal Environ Epidemiol* 14(3):204–213.
- Tan YM, Liao KH, Clewell HJ III. 2007. Reverse dosimetry: interpreting trihalomethanes biomonitoring data using physiologically based pharmacokinetic modeling. *J Expo Sci Environ Epidemiol* 17(7):591–603.
- Tan YM, Liao KH, Conolly RB, Blount BC, Mason AM, Clewell HJ. 2006. Use of a physiologically based pharmacokinetic model to identify exposures consistent with human biomonitoring data for chloroform. *J Toxicol Environ Health A* 69(18):1727–1756.
- U.S. EPA (U.S. Environmental Protection Agency). 2001. Chloroform (CASRN 67-66-3). Available: <http://www.epa.gov/IRIS/subst/0025.htm> [accessed 7 July 2008].
- U.S. EPA (U.S. Environmental Protection Agency). 2007. Drinking Water Contaminants. Available: <http://www.epa.gov/safewater/contaminants/index.html> [accessed 7 July 2008].
- Wallace LA. 1997. Human exposure and body burden for chloroform and other trihalomethanes. *Crit Rev Environ Sci Technol* 27(2):113–194.
- Weisel CP, Little JC, Chiu N, Pandis SN, Davidson C, Wilkes CR. 1999. Developing exposure estimates. In: *Exposure to Contaminants in Drinking Water* (Olin SS, ed). Boca Raton, FL:Chapman and Hall/CRC Press, 85–122.

Table 5. Estimated chloroform intake from drinking water (mg/kg/day).

Percentile	Chloroform intake (mg/kg/day)
5th	9.9×10^{-5}
10th	1.3×10^{-4}
25th	2.1×10^{-4}
50th	4.0×10^{-4}
75th	7.4×10^{-4}
90th	1.2×10^{-3}
95th	1.6×10^{-3}